

**4 month visit**

Today's measurements Wt %, Ht %, HC %

Vaccines

- DTap, IPV, Prevnar, Hib, Rotateq
- Reaction: fever, irritability, redness at site, firm lump at sight. Cool compress today, warm compress tomorrow
- Acetaminophen dosage based on weight- see website for product info and dosing

Nutrition:

- Breast or formula feedings 4-7x/day
- Continue vitamin D for breastfed infants
- Babies may become distracted while feeding, interested in things and people around them. If feeding is difficult, try feeding in a quiet, darkened room.
- Babies start solids between 4-6 months: sign of a baby's readiness are; good head control, interest in watching others eat, opening their mouth as food nears and not sticking tongue out when offer the spoon (extrusion reflex)

Development

- Smiles
- Laughs
- Vocalizes, starting to babble
- Good head control
- Sits with support
- Rolling to side or completely over
- Reaches for objects
- Hands open at rest
- Hands to midline

Teething:

- The first tooth erupts between 4 and 12 months (even up to 18 month in rare instances).
- A cold damp washcloth and chew toys are helpful
- Once teeth erupt, start wiping them with a soft cloth or finger brush and water. No toothpaste is necessary

Sleep

- Develop a sleep/bedtime routine
- Put baby to sleep awake and drowsy
- Baby's can sleep 10-12 hours overnight at this age
- Its ok to start "sleep training"

When to call us:

- Rectal temp >101 for 72 hours
- Crying excessively or inconsolably
- Difficulty breathing
- Lethargic or poor feeding
- Vomiting forcefully or frequently
- OR if you have any other concerns or questions